

Product datasheet for TP728281S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Recombinant R-spondin 1, Human, HEK293

Product data:

Product Type: Recombinant Proteins

Description: Recombinant R-spondin 1, Human, HEK293

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

A DNA sequence encoding Human R-Spondin 1 Protein (#Q2MKA7) (Ser21 - Ala263) was

expressed with polyhistidine-SUMO tag at the N-terminus.

Tag: His-SUMO Tag (N-term)

Predicted MW: The protein has a calculated MW of 46.8 kDa. The protein migrates as 62-68 kDa under

reducing condition (SDS-PAGE analysis).

Purity: >95% as determined by SDS-PAGE.

Buffer: Liquid (1X PBS, pH 7.4)

Bioactivity: R-Spondin-1 enhances BMP-2-mediated differentiation of MC3T3-E1 cells. The ED₅₀ for this

effect is 1-3 µg/mL.

Endotoxin: <1 EU per 1 µg of the protein by the LAL method.

Reconstitution Method: Centrifuge at 3000 rpm for 5 mins before opening. Do Not Vortex! Vigorous shaking may

impair the biological activity of the protein.

Applications: Cell culture

Storage: Store at 2°C to 8°C for up to 2 weeks, -20°C or -80°C for 6 months. Further dilute in a buffer

containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated

freeze/thaw cycles.

UniProt ID: Q2MKA7

Synonyms: RSPO1, Roof plate-specific spondin-1





Summary:

R-spondin 1 (RSPO1, Roof plate-specific Spondin 1) is a cysteine-rich secreted glycoprotein that belongs to the R-Spondin family. Structurally, R-spondin 1 harbors an N-terminal signal peptide, two cysteine-rich furin-like (FU1-FU2) domains, a thrombospondin (TSP) domain, and a basic amino acid-rich (BR) C-terminal domain. R-spondin 1 acts as a ligand for the leucine-rich repeat-containing G-protein coupled receptor (LGR) family (LGR4/5/6), subsequent clearing negative regulators ZNRF3/RNF43 from the membrane, leading to the availability of Wnt receptor and activating Wnt/ β -catenin signaling. The signal cascade is essential for regulating stem cell proliferation, embryonic development, and tissue homeostasis.