

Product datasheet for **TP728286M**

Recombinant TGF beta 2 (Transforming growth factor beta 2), Human

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant TGF beta 2 (Transforming growth factor beta 2), Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MALDAAAYCFRNVQDNCCLRPLYIDFKRDLGWKWIHEPKGYNANFCAGACPYLWSSDTQHSRVLSLYNTI NPEASASPCCVSDLEPLTILYYIGKTPKIEQLSNMIVKSKCS with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 13.66 kDa. The protein migrates as 15 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 3.5.
Bioactivity:	Measure by its ability to inhibit IL-4-induce proliferation in HT-2 cells. The ED ₅₀ for this effect is <0.2 ng/mL. The specific activity of recombinant human TGF beta 2 is > 5 x 10 ⁶ IU/mg.
Endotoxin:	<0.01 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile 10 mM HCl to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	P61812
Synonyms:	G-TSF, LDS4, TGFB2



[View online »](#)

Summary:

Transforming Growth Factors beta 2 (TGF β -2) is a 12.85 kDa member of the epidermal Growth Factors with 113 amino acid residues. TGF β -2 is expressed from throughout the body. TGF β -2 is a regulator of cell proliferation, differentiation, apoptosis, cell plasticity and migration, etc. TGF- β -2 also associates with various kinds of diseases, such as cancer and tissue fibrosis.

Product images:

SDS- PAGE analysis of recombinant human TGF beta 2