

Product datasheet for TP728294

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Recombinant VEGF165 (Vascular endothelial growth factor 165) Human

Product data:

Product Type: Recombinant Proteins

Description: Recombinant VEGF165 (Vascular endothelial growth factor 165) Human

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

MAPMAEGGGQNHHEVVKFMDVYQRSYCHPIETLVDIFQEYPDEIEYIFKPSCVPLMRCGGCCNDEGLECV PTEESNITMQIMRIKPHQGQHIGEMSFLQHNKCECRPKKDRARQENPCGPCSERRKHLFVQDPQTCKCS

CKNTDSRCKARQLELNERTCRCDKPRR with polyhistidine tag at the C-terminus.

Tag: His Tag (C-term)

Predicted MW: The protein has a calculated MW of 20.11 kDa. The protein migrates as 21 kDa under

reducing condition (SDS-PAGE analysis).

Purity: >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 8.0.

Bioactivity: Measure by its ability to induce HUVEC cells proliferation. The ED₅₀ for this effect is <5 ng/mL.

The specific activity of recombinant human VEGF165 is approximately >1.4 x 106 IU/mg.

Endotoxin: $< 0.1 \text{ EU per 1} \mu \text{g of the protein by the LAL method.}$

Reconstitution Method: Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile H_2O to a concentration not less than 100 $\mu g/mL$ and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Applications: Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

RefSeg: NP 001165097

UniProt ID: <u>P15692</u>, <u>A0A0Y0IMM4</u>

Synonyms: VPF, Folliculostellate cell-derived growth factor, Glioma-derived endothelial cell mitogen






Summary:

Vascular Endothelial Growth Factors 165(VEGF165) is a potent growth and angiogenic cytokine which belongs to the VEGF family, includes VEGF-A, VEGF-B, VEGF-C, VEGF-D, VEGF-E, and PIGF. Human VEGF165 is an abundant glycosylated cytokine composed of two identical 165 amino acid chains. Human VEGF165 plays an important role in embryonic vasculogenesis, angiogenesis and neurogenesis.

Product images:



SDS- PAGE analysis of recombinant human VEGF165