

# **Product datasheet for TP728294L**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Recombinant VEGF165 (Vascular endothelial growth factor 165), Human

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant VEGF165 (Vascular endothelial growth factor 165), Human

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

MAPMAEGGGQNHHEVVKFMDVYQRSYCHPIETLVDIFQEYPDEIEYIFKPSCVPLMRCGGCCNDEGLECV PTEESNITMQIMRIKPHQGQHIGEMSFLQHNKCECRPKKDRARQENPCGPCSERRKHLFVQDPQTCKCS

CKNTDSRCKARQLELNERTCRCDKPRR with polyhistidine tag at the C-terminus.

Tag: His Tag (C-term)

**Predicted MW:** The protein has a calculated MW of 20.11 kDa. The protein migrates as 21 kDa under

reducing condition (SDS-PAGE analysis).

**Purity:** >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 8.0.

**Bioactivity:** Measure by its ability to induce HUVEC cells proliferation. The ED<sub>50</sub> for this effect is <5 ng/mL.

The specific activity of recombinant human VEGF165 is approximately >1.4 x 106 IU/mg.

**Endotoxin:**  $< 0.1 \text{ EU per 1} \mu \text{g of the protein by the LAL method.}$ 

**Reconstitution Method:** Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile  $H_2O$  to a concentration not less than 100  $\mu g/mL$  and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

**Applications:** Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

**RefSeg:** NP 001165097

UniProt ID: <u>P15692</u>, <u>A0A0Y0IMM4</u>

**Synonyms:** VPF, Folliculostellate cell-derived growth factor, Glioma-derived endothelial cell mitogen






### **Summary:**

Vascular Endothelial Growth Factors 165(VEGF165) is a potent growth and angiogenic cytokine which belongs to the VEGF family, includes VEGF-A, VEGF-B, VEGF-C, VEGF-D, VEGF-E, and PIGF. Human VEGF165 is an abundant glycosylated cytokine composed of two identical 165 amino acid chains. Human VEGF165 plays an important role in embryonic vasculogenesis, angiogenesis and neurogenesis.

# **Product images:**



SDS- PAGE analysis of recombinant human VEGF165