

Product datasheet for **TP728295**

Recombinant 4-1BBL (4-1BB ligand), Mouse

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant 4-1BBL (4-1BB ligand), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MRTEPRPALTTTSPNLGTRENNADQVTPVSHIGCPNTTQQGSPVFAKLLAKNQASLCNTTLNWHSQDG AGSSYLSQGLRYEEDKKELVVDSPGLYYVFLELKLSPFTNTGCHKVQGWVSLVLQAKPQVDDFDNLALTVE LFPCSMENKLVDRWSQLLLLKAGHRLSVGLRAYLHGAQDAYRDWELSYPNNTTSFGLFLVKPDNPWE with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 23.9 kDa. The protein migrates as 24 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>95% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce IL-2 secretion in mouse T cells in the presence of the anti-CD3 antibody. The ED ₅₀ for this effect is <0.05 µg/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	P41274
Synonyms:	4-1BB, 4-1BB-, 4-1BB-L, 4-1BBL, AI848817, Cd137, Cd137l, Ly6, Ly63l, tumor necrosis factor (ligand) superfamily, member 9 , Tnfsf9



[View online »](#)

Summary:

4-1BB ligand (4-1BBL) is a type II transmembrane protein that is part of the tumor necrosis factor (TNF) ligand family. As an inducible co-stimulatory molecule, it presents on several antigen presenting cell (APC) types, including B cells, macrophages and DCs. The interactions between 4-1BB and 4-1BBL trigger pleiotropic effects on the immune response including antigen presenting process, proliferation of CD4 and CD8 positive T-cells, as well as cytokine secretion from T-cells through NF κ B, c-Jun, and p38 downstream signal pathways activation. Therefore, 4-1BB and 4-1BBL are recently used for the immunotherapy of cancer.

Product images:

SDS- PAGE analysis of recombinant mouse 41BBL