

Product datasheet for **TP728302**

Recombinant CCL4 (C-C Motif Chemokine Ligand 4), Mouse

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant CCL4 (C-C Motif Chemokine Ligand 4), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	APMGSDPPTSCCFSYTSRQLHRSFVMDYYETSSLCSKPAVFLTKRGRQICANPSEPWWTEYMSDLELN with polyhistidine tag at the N-terminus.
Tag:	His Tag (N-term)
Predicted MW:	The protein has a calculated MW of 8.64 kDa. The protein migrates as 11-17 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to chemoattract human PBMCs using a concentration range of 20.0 - 200.0 ng/mL. Note: Results may vary from different PBMC donors.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	P14097
Synonyms:	MIP-1b; Macrophage Inflammatory Protein-1β, ACT-2



[View online »](#)

Summary:

C-C Motif Chemokine Ligand 4 (CCL4) is a 7.66 kDa cytokine with 69 amino acid residues. CCL4, also named macrophage inflammatory protein-1 β (MIP-1 β), is mainly secreted from neutrophils, monocytes, B cells, T cells, fibroblasts, endothelial cells, and epithelial cells. In addition, CCL4 participates in immune responses, including recruitment of immune cells like lymphocytes, monocytes, and leukocytes, response to IL-1 and IFN γ , and production of TNF when CCL4 binds to CCR5.

Product images:

SDS- PAGE analysis of recombinant mouse CCL4