

Product datasheet for **TP728303L**

Recombinant CD27L (CD27 ligand), Mouse

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant CD27L (CD27 ligand), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	QQQRLLLEHPEPHTAELQLNLTVPRKDPTLRWGAGPALGRSFTHGPELEEGHLRIHQDGLYRLHIQVTLAN CSSPGSTLQHRATLAVGICSPAAGHISLLRGRFGDCTVALQRLTYLVHGDVLTCTNLTLPLPSRNADETF GVQWICP with polyhistidine tag at the N-terminus.
Tag:	His Tag (N-term)
Predicted MW:	The protein has a calculated MW of 17.25 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce proliferation in mouse T cells in the presence of the anti-CD3 antibody. The ED ₅₀ for this effect is <4.5 µg/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	O55237
Synonyms:	soluble CD27 Ligand, sCD27 Ligand, TNFSF7, CD70, Tnfs, Tnlg8a



[View online »](#)

Summary:

CD27L is also named for CD70 which is one member of TNF family. CD27L is expressed on T and B lymphocytes and mature DCs. It binds the CD27, which is on the antigen-presenting cells. CD27L is a 19.2 kDa cytokine with 148 amino acid residues which is a transmembrane glycoprotein. CD27L plays an important role in the regulation of T cell proliferation which is also a good target of cancer immunotherapy.

Product images:

SDS- PAGE analysis of recombinant mouse CD27L