

Product datasheet for **TP728319M**

Recombinant FGF-21 (Fibroblast growth factor-21), Mouse

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant FGF-21 (Fibroblast growth factor-21), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	AYPIPDSSPLLQFGGQVRQRYLYTDDDQDTEAHLEIREDGTVGAAHRSPESLLELKALKPGVIQILGVKAS RFLCQQPDGALYGSPHFDPEACSFRELLLEDGYNVYQSEAHGLPLRLPQKDSPNQDATSWGVPVRFPLPMP GLLHEPQDQAGFLPPEPPDVGSSDPLSMVEPLQGRSPSYAS with polyhistidine tag at the N-terminus
Tag:	His Tag (N-term)
Predicted MW:	The protein has a calculated MW of 20.76 kDa. The protein migrates as 25-35 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.
Bioactivity:	Measure by its ability to induce proliferation in NIH-3T3 mouse embryonic fibroblast cells in the presence of mouse Klotho beta and heparin. The ED ₅₀ for this effect is < 2 µg/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	Q9JIN1
Synonyms:	Fgf8c



[View online »](#)

Summary:

Fibroblast Growth Factors 21 (FGF-21) is a 23 kDa protein with 210 amino acid residues. Belongs to FGF superfamily, FGF-21 is related to glucose and lipid metabolism. Peroxisome proliferator-activated receptor gamma (PPAR γ) and peroxisome proliferator-activated receptor alpha (PPAR α) regulate FGF-21 expression.

Product images:

SDS- PAGE analysis of recombinant mouse FGF-21