


Product datasheet for **TP728320**

Recombinant Flt-3 Ligand (Fms-related tyrosine kinase-3 ligand) Mouse

Product data:


| | |
|--|--|
| Product Type: | Recombinant Proteins |
| Description: | Recombinant Flt-3 Ligand (Fms-related tyrosine kinase-3 ligand)  Mouse |
| Species: | Mouse |
| Expression Host: | E. coli |
| Expression cDNA Clone or AA Sequence: | MGTPDCYFSHSPISSNFKVKFRELTDHLLKDYPVTVAVNQLQDEKHCKALWSLFLAQRWIEQLKTAVAGSKM QTLLEDVNTEIHVTSCTFQPLPECLRFVQTNISHLLKDTCTQLLALKPCIGKACQNF SRCLEVQCQPDSST LLPPRSPIALEATELPEPRPR with polyhistidine tag at the C-terminus. |
| Tag: | His Tag (C-term) |
| Predicted MW: | The protein has a calculated MW of 19.3 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis). |
| Purity: | >98% as determined by SDS-PAGE. |
| Buffer: | The protein was lyophilized from a 0.2 μ m filtered solution containing 1X PBS, pH 7.4. |
| Bioactivity: | Measure by its ability to induce proliferation in BaF3 cells transfected with mouse Flt-3. The ED ₅₀ for this effect is <2 ng/mL. |
| Endotoxin: | <0.1 EU per 1 μ g of the protein by the LAL method. |
| Reconstitution Method: | Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 μ g/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein. |
| Applications: | Cell culture |
| Storage: | Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles. |
| UniProt ID: | P49772 |
| Synonyms: | Fetal liver kinase 2, FLK-2 |



[View online »](#)

Summary:

Fms-related tyrosine kinase-3 ligand (Flt-3 Ligand) is a protein which is encoded by the Flt3lg gene in mouse. As an important regulator of hematopoiesis, Flt-3 ligand has a tyrosine-protein kinase activity that stimulate the proliferation of hematopoietic progenitor cells of both lymphoid and myeloid origin. Flt-3 ligand can synergistically increase the proliferation of immature progenitors with other cytokines such as GM-CSF, G-CSF, IL-3, EPO, IL-11, IL-12, IL-6 or TPO. Flt-3 ligand is expressed by a variety of hematopoietic progenitor cells including immature hematopoietic and marrow stromal cells.

Product images:

SDS- PAGE analysis of recombinant mouse Flt-3 Ligand