

Product datasheet for **TP728341S**

Recombinant IL-19 (Interleukin-19), Mouse

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IL-19 (Interleukin-19), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MLRRCLISVDMRLIEKSFHEIKRAMQTKDTFKNVTILSLENLRSIKPGDVCCMTNLLTFYRDRVFQDHQE RSLEVLRRISSIANSFLCVQKSLERCQVHRQCNCSEQATNATRIIHDNYNQLEVSSAALKSLGELNILLAWID RNHLETPAA with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 18.54 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce proliferation in BaF3 cells transfected with human IL-20 R alpha and human IL-20 R beta. The ED ₅₀ for this effect is <0.6 ng/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	Q8CJ70
Synonyms:	Il19



[View online »](#)

Summary:

Interleukin 19 (IL-19) predicts a molecular mass of 35.8 kDa, is a type of anti-inflammatory cytokine. It promotes the Th2 T-cell response which supports an anti-inflammatory lymphocyte phenotype, dampens the Th1 T-cell response and IFN γ secretion, increases IL-10 expression in peripheral blood mononuclear cells, and inhibits the production of IgG from B cells.

Product images:

SDS- PAGE analysis of recombinant mouse IL-19