

Product datasheet for **TP728342L**

Recombinant IL-1RA (Interleukin-1 receptor antagonist), Mouse

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IL-1RA (Interleukin-1 receptor antagonist), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MRPSGKRPCMKQAFRIWDTNQKTFYLRNNQLIAGYLQGPNIKLEEKIDMVPIDLHSVFLGIHGGKLCCLSC AKSGDDIKLQLEEVNITDLSKNKEEDKRFTFIRSEKGPSTSFESAACPGWFLCTTLEADRPVSLTNTPEEPLIV TKFYFQEDQ with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 18.27 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to inhibit IL-1 alpha -dependent proliferation in D10.G4.1 cells. The ED ₅₀ for this effect is <50 ng/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	P25085
Synonyms:	ICIL-1RA, IRAP, IL-1RN, F630041P17Rik



[View online »](#)

Summary:

Interleukin 1 receptor antagonist (IL-1RA) predicts a molecular mass of 19.8 kDa, is a member of the IL-1 family that binds to IL-1 receptors but does not induce any intracellular response. Two structural variants of IL-1RA have previously been described: a 17 kDa form that is secreted from monocytes, macrophages, neutrophils, and other cells (sIL-1RA) and an 18 kDa form that remains in the cytoplasm of keratinocytes and other epithelial cells, monocytes, and fibroblasts (icIL-1RA).

Product images:

SDS- PAGE analysis of recombinant mouse IL-1RA