

## Product datasheet for **TP728347**

### Recombinant IL-24 (Interleukin-24), Mouse

#### Product data:


|  |   |
|--|---|
| <b>Product Type:</b>                         | Recombinant Proteins  |
| <b>Description:</b>                          | Recombinant IL-24 (Interleukin-24), Mouse   |
| <b>Species:</b>                              | Mouse   |
| <b>Expression Host:</b>                      | E. coli   |
| <b>Expression cDNA Clone or AA Sequence:</b> | MQEFRFGSCQVTGWLPPELWEAFWTVKNTVQTQDDITSIRLLKPQVLRNVSGAESCYLHSLKLFYLNVTFKNYHSKIAKFKVLRFSSTLANNFIVIMSQLQPSKDNSMLPISESAHQRFLLFRRAFKQLDTEVALVKAFGEVDILLTWMQKFYHL with polyhistidine tag at the C-terminus.   |
| <b>Tag:</b>                                  | His Tag (C-term)  |
| <b>Predicted MW:</b>                         | The protein has a calculated MW of 18.94 kDa. The protein migrates about 17 kDa under reducing condition (SDS-PAGE analysis).   |
| <b>Purity:</b>                               | >95% as determined by SDS-PAGE.   |
| <b>Buffer:</b>                               | The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.  |
| <b>Bioactivity:</b>                          | Measure by its ability to induce proliferation in BaF3 cells transfected with human IL-20 R alpha and human IL-20 R beta. The ED <sub>50</sub> for this effect is <0.3 ng/mL.   |
| <b>Endotoxin:</b>                            | <0.1 EU per 1 µg of the protein by the LAL method.  |
| <b>Reconstitution Method:</b>                | Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein. |
| <b>Applications:</b>                         | Cell culture  |
| <b>Storage:</b>                              | Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.                        |
| <b>RefSeq:</b>                               | <a href="#">NP_444325.3</a>   |
| <b>Synonyms:</b>                             | MDA-7 (Melanoma Differentiation-Associated gene 7 protein), FISP, St16  |



[View online »](#)

**Summary:**

Interleukin 24 (IL-24) predicts a molecular mass of 18.2 kDa, is a cytokine belonging to the IL-10 family of cytokines that signals through IL-20R1 /IL-20R2 and IL-22R1 /IL-20R2. Its normal physiological role is connected with wound healing, protection against diseases caused by bacteria.

**Product images:**

SDS- PAGE analysis of recombinant mouse IL-24