

Product datasheet for TP728347L

Recombinant IL-24 (Interleukin-24), Mouse

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IL-24 (Interleukin-24), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MQEFRFGSCQVTGWLPPELWEAFWTVKNTVQTQDDITSIRLLKPQVLRNVSGAESCYLHSLKLFYLNVTFKNYHSKIAKFKVLRFSSTLANNFIVIMSQLQPSKDNSMLPISESAHQRFLLFRRAFKQLDTEVALVKAFGEVDILLTWMQKFYHL with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 18.94 kDa. The protein migrates about 17 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>95% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce proliferation in BaF3 cells transfected with human IL-20 R alpha and human IL-20 R beta. The ED ₅₀ for this effect is <0.3 ng/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
RefSeq:	NP_444325.3
Synonyms:	MDA-7 (Melanoma Differentiation-Associated gene 7 protein), FISP, St16



[View online »](#)

Summary:

Interleukin 24 (IL-24) predicts a molecular mass of 18.2 kDa, is a cytokine belonging to the IL-10 family of cytokines that signals through IL-20R1 /IL-20R2 and IL-22R1 /IL-20R2. Its normal physiological role is connected with wound healing, protection against diseases caused by bacteria.

Product images:

SDS- PAGE analysis of recombinant mouse IL-24