

Product datasheet for TP728348

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Recombinant IL-25 (Interleukin-25), Mouse

Product data:

Product Type: Recombinant Proteins

Description: Recombinant IL-25 (Interleukin-25), Mouse

Species: Mouse Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

MVSLRIQEGCSHLPSCCPSKEQEPPEEWLKWSSASVSPPEPLSHTHHAESCRASKDGPLNSRAISPWSYEL DRDLNRVPQDLYHARCLCPHCVSLQTGSHMDPLGNSVPLYHNQTVFYRRPCHGEEGTHRRYCLERRLYR

VSLACVCVRPRVMA with polyhistidine tag at the C-terminus.

Tag: His Tag (C-term)

Predicted MW: The protein has a calculated MW of 18.41kDa. The protein migrates about 17 kDa under

reducing condition (SDS-PAGE analysis).

Purity: >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate,

0.2 M NaCl, pH 4.5.

Bioactivity: Measured by its ability to induce CXCL1 secretion in HT-29 cells. The ED₅₀ for this effect is <1

ng/mL

Endotoxin: <0.1 EU per 1 μg of the protein by the LAL method.

Reconstitution Method: Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile H_2O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Applications: Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

 RefSeq:
 NP 542767 1

 Synonyms:
 IL-17E, II17e






Summary:

Interleukin 25 (IL-25) is a cytokine with proinflammatory and anti-inflammatory effects, it predicts a molecular mass of 16.7 kDa. It can induce NF-κB activation, and stimulate the production of IL-8, which is the major chemotactic substance of neutrophils.

Product images:



SDS- PAGE analysis of recombinant mouse IL-25