

## **Product datasheet for TP728348M**

## OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## Recombinant IL-25 (Interleukin-25), Mouse

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant IL-25 (Interleukin-25), Mouse

Species: Mouse Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

MVSLRIQEGCSHLPSCCPSKEQEPPEEWLKWSSASVSPPEPLSHTHHAESCRASKDGPLNSRAISPWSYEL DRDLNRVPQDLYHARCLCPHCVSLQTGSHMDPLGNSVPLYHNQTVFYRRPCHGEEGTHRRYCLERRLYR

VSLACVCVRPRVMA with polyhistidine tag at the C-terminus.

Tag: His Tag (C-term)

**Predicted MW:** The protein has a calculated MW of 18.41kDa. The protein migrates about 17 kDa under

reducing condition (SDS-PAGE analysis).

**Purity:** >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate,

0.2 M NaCl, pH 4.5.

**Bioactivity:** Measured by its ability to induce CXCL1 secretion in HT-29 cells. The ED<sub>50</sub> for this effect is <1

ng/mL

**Endotoxin:** <0.1 EU per 1 μg of the protein by the LAL method.

**Reconstitution Method:** Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile  $H_2O$  to a concentration not less than 100  $\mu$ g/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

**Applications:** Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

 RefSeq:
 NP 542767 1

 Synonyms:
 IL-17E, II17e






**Summary:** 

Interleukin 25 (IL-25) is a cytokine with proinflammatory and anti-inflammatory effects, it predicts a molecular mass of 16.7 kDa. It can induce NF-κB activation, and stimulate the production of IL-8, which is the major chemotactic substance of neutrophils.

## **Product images:**



SDS- PAGE analysis of recombinant mouse IL-25