

## Product datasheet for **TP728352L**

### Recombinant IL-3 (Interleukin-3), Mouse


#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant IL-3 (Interleukin-3), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MDTHRLTRLNCSSIVKEIIGKLPPELKTDDGPSLRNKSFRRVNLSKFVESQGEVDPEDRYVIKSNLQKLN CCLPTSANDSALPGVFIRDLDLDFRKKLRFYMHVHLNDLETVLTSRPPQPASGSVSPNRGTVEC with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 16.04 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce NFS-60 cells proliferation. The ED <sub>50</sub> for this effect is <85 pg/mL. The specific activity of recombinant mouse IL-3 is approximately >1 x 10 <sup>7</sup> IU/mg.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	<a href="#">P01586</a>
Synonyms:	MCGF (Mast Cell Growth Factor), Multi-CSF, HCGF, P-cell stimulation factor, Interleukin-3b, BPA, PSF, Csfmu


[View online »](#)

**Summary:**

Interleukin-3 (IL-3) is a pleiotropic cytokine which can stimulates the survival, differentiation and proliferation of committed progenitor cells, including megakaryocyte, granulocyte-macrophage, erythroid, eosinophil, basophil and mast cell lineages. IL-3 also enhances phagocytosis and antibody-mediated cellular cytotoxicity. IL-3 binds to IL-3R (IL-3 receptor), which is composed of a unique  $\alpha$  subunit (IL-3R $\alpha$ ) and a common  $\beta$ -subunit ( $\beta$ c).

**Product images:**


SDS- PAGE analysis of recombinant mouse IL-3