

Product datasheet for **TP728354S**

Recombinant IL-33 (Interleukin-33), Mouse

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IL-33 (Interleukin-33), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MSIQGTSLLTQSPASLSTYNDQSVSFVLENGCYVINVDDSGKDQEQDQVLLRRYYESPCPASQSGDGVGDKLMLVMNSPIKDTDIWLHANDKDYSVELQRGDVSPPEQAFFVLHKKSSDFVSFECKNLPGTYIGVKDNQLALVEEKDESCNNIMFKLSKI with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 18.51 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce proliferation in D10.G4.1 cells. The ED ₅₀ for this effect is <40 pg/mL. The specific activity of recombinant mouse IL-33 is > 2 x 10 ⁷ IU/mg.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	Q8BVZ5
Synonyms:	IL-1 F11, NF-HEV, 9230117N10Rik



[View online »](#)

Summary:

Interleukin 33 (IL-33) is a 17.65 kDa cytokine with 159 amino acid residues. IL-33 is a member of the IL-1 family and secreted from various cell types, such as mast cells, macrophages, endothelial cells, and epithelial cells. IL-33 activates NF- κ B and MAPK signaling pathways that stimulate the secretion of type 2 cytokines like IL-5 and IL-13 when IL-33 binds to the ST2 receptor primarily expressed in mast cells and Th2 cells. In addition to Th type 2 responses, IL-33 participates in maintaining barrier tissue defense.

Product images:

SDS- PAGE analysis of recombinant mouse IL-33