

Product datasheet for **TP728356S**

Recombinant IL-36 alpha (Interleukin-36 alpha), Mouse

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IL-36 alpha (Interleukin-36 alpha), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MNKEKELRAASPSLRHVQDLSSRVWILQNNILTAVPRKEQTVPVTTILLPCQYLDLTLETNRGDPTYMGVQR PMSCLFCTKDGEQPVQLQEGGNIMEMYNKKEPVKASLFYHKKSGTTSTFESAAPGWFIAVCSKGCPLIL TQELGEIFITDFEMIVVH with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 18.82 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED ₅₀ for this effect is <15 ng/mL. The specific activity of recombinant mouse IL-36 alpha is > 1 x 10 ⁵ IU/mg.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	Q9JLA2
Synonyms:	IL36A, Interleukin-1 family member 6 (IL-1F6), FIL1ε (FIL1E), Interleukin-1ε (IL1E), Fi, Fil, Fil1, IL-1, IL-1H1, IL1RP2



[View online »](#)

Summary:

Interleukin 36 alpha (IL-36 alpha) is a 17.76 kDa cytokine with 160 amino acid residues. IL-36 alpha is an IL-1 family member, which binds to the Interleukin-1 receptor and is expressed in the lung, stomach, and placenta. IL-36 alpha is critical in inflammatory responses and stimulates the production of proinflammatory cytokines, such as IL-12, IL-1 β , IL-6, and TNF- α . It also induces the generation of IFN γ , IL-4, and IL-17 in CD4+ T cells.

Product images:

SDS- PAGE analysis of recombinant mouse IL-36 alpha