

## **Product datasheet for TP728360S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Recombinant IL-5 (Interleukin-5), Mouse

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant IL-5 (Interleukin-5), Mouse

Species: Mouse Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

 ${\tt MEIPMSTVVKETLTQLSAHRALLTSNETMRLPVPTHKNHQLCIGEIFQGLDILKNQTVRGGTVEMLFQNLS}$ 

LIKKYIDRQKEKCGEERRRTRQFLDYLQEFLGVMSTEWAMEG with polyhistidine tag at the C-

terminus.

Tag: His Tag (C-term)

Predicted MW: The protein has a calculated MW of 13.93 kDa. The protein migrates as 11-17 kDa under

reducing condition (SDS-PAGE analysis).

**Purity:** >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 7.4.

**Bioactivity:** Measure by its ability to induce TF-1 cells proliferation. The ED<sub>50</sub> for this effect is < 0.2 ng/mL.

The specific activity of recombinant mouse IL-5 is  $> 5 \times 10^6$  IU/mg.

**Endotoxin:**  $< 0.1 \text{ EU per 1} \mu \text{g of the protein by the LAL method.}$ 

**Reconstitution Method:** Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile  $H_2O$  to a concentration not less than 100  $\mu g/mL$  and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

**Applications:** Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

UniProt ID: P04401

Synonyms: EDF, BCDFII, TRF






**Summary:** 

Interleukin 5 (IL-5) is a 45-60 kDa homodimeric cytokine that functions principally in the eosinophil arm of the Th2 response. It is a member of the helical bundle family of cytokines, along with IL-3, IL-4, and granulocyte macrophage-colony stimulating factor (GM-CSF).

## **Product images:**



SDS- PAGE analysis of recombinant mouse IL-5