

## Product datasheet for TP728367M

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Recombinant Neurturin, Mouse**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant Neurturin, Mouse

Species: Mouse Expression Host: E. coli

**Expression cDNA Clone** PGARPCGLRELEVRVSELGLGYTSDETVLFRYCAGACEAAIRIYDLGLRRLRQRRRVRRERARAHPCCRPTAY

or AA Sequence: EDEVSFLDVHSRYHTLQELSARECACV with polyhistidine tag at the N-terminus.

Tag: His Tag (N-term)

Predicted MW: The protein has a calculated MW of 12.33 kDa. The protein migrates as 11-17 kDa under

reducing condition (SDS-PAGE analysis).

**Purity:** >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 7.4.

**Bioactivity:** Measure by its ability to induce proliferation in SH-SY5Y cells. The ED₅₀ for this effect is <50

ng/mL.

**Endotoxin:**  $< 0.1 \; EU \; per \; 1 \; \mu g \; of \; the \; protein \; by \; the \; LAL \; method.$ 

**Reconstitution Method:** Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile  $H_2O$  to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

**Applications:** Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

UniProt ID: P97463

Synonyms: NTN, NRTN






**Summary:** 

Neurturin is a 22 kDa cytokine with 195 amino acid residues. Neurturin belongs to the same neurotrophic factor family as GDNF. Binding to GFR $\alpha$ -2 (GPI-linked receptor termed GDNF family receptor  $\alpha$ -2), neurturin activates Ret receptor tyrosine kinase and MAPK pathway. It involves in neuron survival.

## **Product images:**



SDS- PAGE analysis of recombinant mouse Neurturin