

Product datasheet for TP728368

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Recombinant Noggin, Mouse, HEK293

Product data:

Product Type: Recombinant Proteins

Description: Recombinant Noggin, Mouse, HEK293

Species: Mouse Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

A DNA sequence encoding Mouse Noggin(#P97466) (Met1-Cys232) was expressed with

Human IgG1 Fc tag at the C-terminus.

Tag: Human IgG1 Fc Tag (C-term)

Predicted MW: The protein has a calculated MW of 49.14 kDa. The protein migrates as 50 kDa under

reducing condition (SDS-PAGE analysis).

Purity: >95% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 7.4.

Bioactivity: Testing in process

Endotoxin: <0.1 EU per 1 µg of the protein by the LAL method.

Reconstitution Method: Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile H_2O to a concentration not less than 100 μ g/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Applications: Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

UniProt ID: P97466

Synonyms: Nog






Summary:

Noggin binds members of the transforming Growth Factors-beta (TGF beta) superfamily signaling proteins, such as bone morphogenetic protein-4 (BMP-4), which inactivates their activities. As a extracellular antagonist of BMP proteins, Noggin involves in the development of many body tissues, including nerve tissue, muscles, and bones. In addition, Noggin is able to inhibit chondrocyte differentiation through its interaction with GDF5.

Product images:



SDS- PAGE analysis of recombinant mouse Noggin