

# **Product datasheet for TP728374M**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Recombinant TRAIL (TNF-related apoptosis-inducing ligand), Mouse

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant TRAIL (TNF-related apoptosis-inducing ligand), Mouse

Species: Mouse Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

MPRGGRPQKVAAHITGITRRSNSALIPISKDGKTLGQKIESWESSRKGHSFLNHVLFRNGELVIEQEGLYYIY SQTYFRFQEAEDASKMVSKDKVRTKQLVQYIYKYTSYPDPIVLMKSARNSCWSRDAEYGLYSIYQGGLFELK

KNDRIFVSVTNEHLMDLDQEASFFGAFLIN with polyhistidine tag at the C-terminus.

Tag: His Tag (C-term)

Predicted MW: The protein has a calculated MW of 21.00 kDa. The protein migrates as 17-25 kDa under

reducing condition (SDS-PAGE analysis).

**Purity:** >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 7.4.

**Bioactivity:** Measure by its ability to induce cytotoxicity in L929 cells in the presence of actinomycin D.

The ED<sub>50</sub> for this effect is <1 ng/mL.

**Endotoxin:** <0.1 EU per 1 µg of the protein by the LAL method.

**Reconstitution Method:** Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile  $H_2O$  to a concentration not less than 100  $\mu g/mL$  and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

**Applications:** Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

UniProt ID: P50592

Synonyms: TNFSF10, Apo2 Ligand, TL2, Apo2L, CD253






#### **Summary:**

Tumor Necrosis Factor associated Apoptosis Inducing Ligand (TRAIL) is a type II membrane protein of the tumor necrosis factor (TNF) family members, moreover expressed in many adult tissues including the thymus, prostate, colon, ovary and lung. TRAIL is a 19 kDa protein containing 281 residues. Mouse TRAIL shares 70% amino acids sequence identity with human, bovine, and porcine. TRAIL to induce apoptosis in human breast carcinoma cells (MCF7) and human epithelioid carcinoma (HeLa) cell lines, by activate two death receptors of DR4 and DR5 or two decoy receptors DcR1 and DcR2.

### **Product images:**



SDS- PAGE analysis of recombinant mouse TRAIL