

Product datasheet for **TP728375L**

Recombinant TWEAK (TNF-related weak inducer of apoptosis), Mouse


Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant TWEAK (TNF-related weak inducer of apoptosis), Mouse
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MSAPKGRKARPRRAIAAHYEVHPRPGDGAQAGVDGTVSGWEETKINSSSPLRYDRQIGFTVIRAGLYYL YCQVHFDEGKAVYLKDLLVNGVLALRCLEEFSAASSPGPQLRLCQVSGLLPLRPGSSLRIRTLPPWAHLK AAPFLTYFGLFQVH with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 18.14 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce proliferation in HUVEC cells. The ED ₅₀ for this effect is <0.2 µg/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	O54907
Synonyms:	TNFSF12, DR3LG, Apo3 Ligand, AI255180, C87282, Fn1, Fn14, HPIP, TWEAK-R, Twea, TweakR, Tnfrsf12a


[View online »](#)

Summary:

TNF-related Weak Inducer of Apoptosis (TWEAK) is a type II membrane protein of the tumor necrosis factor (TNF) family members, high levels expression in mouse peritoneal macrophages. The murine and human TWEAK homologs are very close related (~93% sequence identity in the receptor-binding domain). TWEAK is a 17.3 kDa protein containing 129 residues. TWEAK induce apoptosis through cognate receptor Fn14 activate caspase-8 and caspase-3, in HSC3 cells KATO-III gastric adenocarcinoma cells and IFN- γ -treated HT-29. TWEAK stimulate inflammatory cytokines in HT-29, A375 cells, WI-38 fibroblast and astrocytes.

Product images:


SDS- PAGE analysis of recombinant mouse TWEAK