

Product datasheet for TP750136

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TRIM72 (NM 001008274) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human MG53, Ala58-End, tag free, expressed in E.coli, 50ug

Species: Human E. coli **Expression Host:**

Expression cDNA Clone

or AA Sequence:

A DNA sequence encoding the region(Ala58-End) of MG53

Tag: Tag Free Predicted MW: 46.9 kDa

>0.05 µg/µL as determined by microplate BCA method **Concentration:**

Purity: > 90% as determined by SDS-PAGE and Coomassie blue staining

Buffer: PBS, pH 7.4, 10% glycerol

Endotoxin: < 1 EU per 1 µg of the protein by the LAL

For testing in cell culture applications, please filter before use. Note that you may experience Note:

some loss of protein during the filtration process.

Store at -80°C. Storage:

Stable for 12 months from the date of receipt of the product under proper storage and Stability:

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001008275

Locus ID: 493829

UniProt ID: Q6ZMU5, A0A590UJ49

RefSeq Size: 2098

Cytogenetics: 16p11.2

RefSeq ORF: 1431

Synonyms: MG53






Summary:

Muscle-specific protein that plays a central role in cell membrane repair by nucleating the assembly of the repair machinery at injury sites. Specifically binds phosphatidylserine. Acts as a sensor of oxidation: upon membrane damage, entry of extracellular oxidative environment results in disulfide bond formation and homooligomerization at the injury site. This oligomerization acts as a nucleation site for recruitment of TRIM72-containing vesicles to the injury site, leading to membrane patch formation. Probably acts upstream of the Ca(2+)-dependent membrane resealing process. Required for transport of DYSF to sites of cell injury during repair patch formation. Regulates membrane budding and exocytosis. May be involved in the regulation of the mobility of KCNB1-containing endocytic vesicles (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Purified recombinant protein MG53 was analyzed by SDS-PAGE gel and Coomossie Blue Staining.