

Product datasheet for TP750194-B

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

KRAS (NM 004985) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human v-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog

(KRAS), transcript variant b, full length, with N-terminal AVI tag, expressed in E.coli, 50ug

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

A DNA sequence encoding the full-length of KRAS

Tag: N-AVI

Predicted MW: 21.2 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by Bradford protein assay method. Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 50 mM Tris-HCl, pH 8.0, 500 mM NaCl, 10% glycerol

Bioactivity: The biotin to protein ratio is 1 as determined by Streptavidin Pull-Down Assay.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 004976

 Locus ID:
 3845

 UniProt ID:
 P01116

 RefSeq Size:
 5312

 Cytogenetics:
 12p12.1

 RefSeq ORF:
 564

Synonyms: C-K-RAS; c-Ki-ras2; CFC2; K-Ras; K-RAS2A; K-RAS2B; K-RAS4B; KI-RAS; KRAS1; KRAS2;

NS; NS3; RALD; RASK2





KRAS (NM_004985) Human Recombinant Protein - TP750194-B

Summary:

This gene, a Kirsten ras oncogene homolog from the mammalian ras gene family, encodes a protein that is a member of the small GTPase superfamily. A single amino acid substitution is responsible for an activating mutation. The transforming protein that results is implicated in various malignancies, including lung adenocarcinoma, mucinous adenoma, ductal carcinoma of the pancreas and colorectal carcinoma. Alternative splicing leads to variants encoding two isoforms that differ in the C-terminal region. [provided by RefSeq, Jul 2008]

Protein Families:

Druggable Genome

Protein Pathways:

Acute myeloid leukemia, Axon guidance, B cell receptor signaling pathway, Bladder cancer, Chemokine signaling pathway, Chronic myeloid leukemia, Colorectal cancer, Dorso-ventral axis formation, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Gap junction, Glioma, GnRH signaling pathway, Insulin signaling pathway, Long-term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Melanoma, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Non-small cell lung cancer, Pancreatic cancer, Pathways in cancer, Progesterone-mediated oocyte maturation, Prostate cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, T cell receptor signaling pathway, Thyroid cancer, Tight junction, VEGF signaling pathway