

Product datasheet for TP761109

UCP2 (NM_003355) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins Description: Purified recombinant protein of Human uncoupling protein 2 (mitochondrial, proton carrier) (UCP2), nuclear gene encoding mitochondrial protein, full length, with N-terminal HIS tag, expressed in E. coli, 50ug Species: Human **Expression Host:** F. coli **Expression cDNA Clone** A DNA sequence encoding human full-length UCP2 or AA Sequence: Tag: N-His Predicted MW: 33 kDa **Concentration:** >0.05 µg/µL as determined by microplate BCA method > 80% as determined by SDS-PAGE and Coomassie blue staining **Purity: Buffer:** 25 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1% sarkosyl, 10% glycerol For testing in cell culture applications, please filter before use. Note that you may experience Note: some loss of protein during the filtration process. Store at -80°C. Storage: Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. NP 003346 RefSeq: 7351 Locus ID: UniProt ID: P55851 **RefSeq Size:** 1646 Cytogenetics: 11q13.4 **RefSeq ORF:** 927 Synonyms: BMIQ4; SLC25A8; UCPH



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRIGENE UCP2 (NM_003355) Human Recombinant Protein – TP761109

Summary: Mitochondrial uncoupling proteins (UCP) are members of the larger family of mitochondrial anion carrier proteins (MACP). UCPs separate oxidative phosphorylation from ATP synthesis with energy dissipated as heat, also referred to as the mitochondrial proton leak. UCPs facilitate the transfer of anions from the inner to the outer mitochondrial membrane and the return transfer of protons from the outer to the inner mitochondrial membrane. They also reduce the mitochondrial membrane potential in mammalian cells. Tissue specificity occurs for the different UCPs and the exact methods of how UCPs transfer H+/OH- are not known. UCPs contain the three homologous protein domains of MACPs. This gene is expressed in many tissues, with the greatest expression in skeletal muscle. It is thought to play a role in nonshivering thermogenesis, obesity and diabetes. Chromosomal order is 5'-UCP3-UCP2-3'. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome

Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US