

Product datasheet for TP761118

MAX (NM_002382) Human Recombinant Protein

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Recombinant Proteins	
Description:	Purified recombinant protein of Human MYC associated factor X (MAX), transcript variant 1, full length, with N-terminal HIS tag, expressed in E. coli, 50ug	
Species:	Human	
Expression Host:	E. coli	
Expression cDNA Clone or AA Sequence:	A DNA sequence encoding human full-length MAX	
Tag:	N-His	
Predicted MW:	18.1 kDa	
Concentration:	>0.05 µg/µL as determined by microplate BCA method	
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining	
Buffer:	25 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1% sarkosyl, 10% glycerol	
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.	
Storage:	Store at -80°C.	
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.	
RefSeq:	<u>NP 002373</u>	
Locus ID:	4149	
UniProt ID:	<u>P61244, Q8TAX8</u>	
RefSeq Size:	2068	
Cytogenetics:	14q23.3	
RefSeq ORF:	480	
Synonyms:	bHLHd4	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	MAX (NM_002382) Human Recombinant Protein – TP761118
Summary:	The protein encoded by this gene is a member of the basic helix-loop-helix leucine zipper (bHLHZ) family of transcription factors. It is able to form homodimers and heterodimers with other family members, which include Mad, Mxi1 and Myc. Myc is an oncoprotein implicated in cell proliferation, differentiation and apoptosis. The homodimers and heterodimers compete for a common DNA target site (the E box) and rearrangement among these dimer forms provides a complex system of transcriptional regulation. Mutations of this gene have been reported to be associated with hereditary pheochromocytoma. A pseudogene of this gene is located on the long arm of chromosome 7. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2012]
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathway	MAPK signaling pathway, Pathways in cancer, Small cell lung cancer

Product images:

116 -	-
66 -	-
45 -	_
35 -	-
25 -	
18 -	_
14 -	-

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US