

Product datasheet for TP761588

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Endothelin 3 (EDN3) (NM_000114) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human endothelin 3 (EDN3), transcript variant 1, full length,

with N-terminal GST and C-terminal HIS tag, expressed in E. coli, 50ug

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

A DNA sequence encoding human full-length EDN3

Tag: N-GST and C-His

Predicted MW: 50.8 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1% sarkosyl, 10% glycerol

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 000105

 Locus ID:
 1908

 UniProt ID:
 P14138

 RefSeq Size:
 2438

 Cytogenetics:
 20q13.32

RefSeq ORF: 714

Synonyms: ET-3; ET3; HSCR4; PPET3; WS4B





Summary:

The protein encoded by this gene is a member of the endothelin family. Endothelins are endothelium-derived vasoactive peptides involved in a variety of biological functions. The active form of this protein is a 21 amino acid peptide processed from the precursor protein. The active peptide is a ligand for endothelin receptor type B (EDNRB). The interaction of this endothelin with EDNRB is essential for development of neural crest-derived cell lineages, such as melanocytes and enteric neurons. Mutations in this gene and EDNRB have been associated with Hirschsprung disease (HSCR) and Waardenburg syndrome (WS), which are congenital disorders involving neural crest-derived cells. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Oct 2014]

Protein Families:

Druggable Genome, Secreted Protein

Product images:

