

Product datasheet for TP761779

KLRC4 (NM_013431) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins Description: Purified recombinant protein of Human killer cell lectin-like receptor subfamily C, member 4 (KLRC4), full length, with N-terminal GST and C-terminal His tag, expressed in E. coli, 50ug Species: Human **Expression Host:** E. coli **Expression cDNA Clone** A DNA sequence encoding human full-length KLRC4 or AA Sequence: N-GST and C-His Tag: Predicted MW: 46.1 kDa **Concentration:** >0.05 µg/µL as determined by microplate BCA method **Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** 50 mM Tris-HCl, pH 8.0, 8 M urea Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. Store at -80°C. Storage: Stable for 12 months from the date of receipt of the product under proper storage and Stability: handling conditions. Avoid repeated freeze-thaw cycles. **RefSeq:** NP 038459 Locus ID: 8302 **UniProt ID:** 043908 **RefSeq Size:** 928 Cytogenetics: 12p13.2 **RefSeq ORF:** 474 Synonyms: NKG2-F; NKG2F



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

KLRC4 (NM_013431) Human Recombinant Protein - TP761779

Summary: Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. This gene is a member of the NKG2 group of genes that are expressed primarily in natural killer (NK) cells. These family members encode transmembrane proteins that are characterized by a type II membrane orientation (have an extracellular C-terminus) and the presence of a C-type lectin domain. This family member is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. Read-through transcription exists between this gene and the downstream KLRK1 (killer cell lectin-like receptor subfamily K, member 1) family member. [provided by RefSeq, Dec 2010]

Protein Families: Transmembrane

Protein Pathways:

Antigen processing and presentation

Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US