

Product datasheet for TP762398

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD45 (PTPRC) (NM_002838) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human protein tyrosine phosphatase, receptor type, C

(PTPRC), transcript variant 1, Thr221-Gln396, with N-terminal His tag, expressed in E.coli, 50ug

Species: Human Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

A DNA sequence encoding the region (Thr221-Gln396,) of CD45

Tag: N-His

Predicted MW: 20.0 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: >80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 50 mM Tris-HCl, pH 8.0, 8 M urea

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 002829

 Locus ID:
 5788

 UniProt ID:
 P08575

 RefSeq Size:
 5026

Cytogenetics: 1q31.3-q32.1

RefSeq ORF: 3912

Synonyms: B220; CD45; CD45R; GP180; L-CA; LCA; LY5; T200






Summary:

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq, Jun 2012]

Protein Families: Protein Pathways: Druggable Genome, ES Cell Differentiation/IPS, Phosphatase, Transmembrane Cell adhesion molecules (CAMs), Fc gamma R-mediated phagocytosis, Primary immunodeficiency, T cell receptor signaling pathway

Product images:



Purified recombinant protein CD45 was analyzed by SDS-PAGE gel and Coomossie Blue Staining.