

# Product datasheet for TP762403

### c-Myc (MYC) (NM\_002467) Human Recombinant Protein

#### **Product data:**

Product Type:	Recombinant Proteins		
Description:	Purified recombinant protein of Human v-myc myelocytomatosis viral oncogene homolog (avian) (Myc), Leu1-Asp343, with N-terminal His tag, expressed in E.coli, 50ug		
Species:	Human		
Expression Host:	E. coli		
Expression cDNA Clone or AA Sequence:	A DNA sequence encoding the region(Leu1-Asp343) of c-Myc		
Tag:	N-His		
Predicted MW:	37.4 kDa		
Concentration:	>0.05 µg/µL as determined by microplate BCA method		
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining		
Buffer:	50 mM Tris-HCl, pH 8.0, 8 M urea		
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.		
Storage:	Store at -80°C after receiving vials.		
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.		
RefSeq:	<u>NP 002458</u>		
Locus ID:	4609		
UniProt ID:	<u>P01106</u>		
RefSeq Size:	2379		
Cytogenetics:	8q24.21		
RefSeq ORF:	1362		
Synonyms:	bHLHe39; c-Myc; MRTL; MYCC		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	c-Myc (MYC) (NM_002467) Human Recombinant Protein – TP762403	
This gene is a proto-oncogene and encodes a nuclear phosphoprotein that place cycle progression, apoptosis and cellular transformation. The encoded protein heterodimer with the related transcription factor MAX. This complex binds to t consensus sequence and regulates the transcription of specific target genes. A this gene is frequently observed in numerous human cancers. Translocations i gene are associated with Burkitt lymphoma and multiple myeloma in human p is evidence to show that translation initiates both from an upstream, in-frame and a downstream AUG start site, resulting in the production of two isoforms termini. [provided by RefSeq, Aug 2017]		
Protein Families:	Druggable Genome, Embryonic stem cells, Induced pluripotent stem cells, Stem cell - Pluripotency, Stem cell relevant signaling - JAK/STAT signaling pathway, Stem cell relevant signaling - TGFb/BMP signaling pathway, Stem cell relevant signaling - Wnt Signaling pathway, Transcription Factors	
Protein Pathways	Acute myeloid leukemia, Bladder cancer, Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, ErbB signaling pathway, Jak-STAT signaling pathway, MAPK signaling pathway, Pathways in cancer, Small cell lung cancer, TGF-beta signaling pathway, Thyroid cancer, Wnt signaling pathway	

## Product images:

116 —	-
66 —	-
45 —	-
35 —	-
25 —	-
18	
14 —	•

Purified recombinant protein c-Myc was analyzed by SDS-PAGE gel and Coomossie Blue Staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US