

Product datasheet for TP762409

MET (NM_000245) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins Description: Purified recombinant protein of Human met proto-oncogene (hepatocyte growth factor receptor) (MET), transcript variant 2, Phe146-Thr230-GGGGS-Ile850-Leu920, with N-terminal His tag, expressed in E.coli, 50ug Species: Human **Expression Host:** F. coli **Expression cDNA Clone** A DNA sequence encoding the region (Phe146-Thr230-GGGGS-Ile850-Leu920) of MET or AA Sequence: Tag: N-His Predicted MW: 17.6 kDa **Concentration:** >0.05 µg/µL as determined by microplate BCA method >80% as determined by SDS-PAGE and Coomassie blue staining **Purity: Buffer:** 50 mM Tris-HCl, pH 8.0, 8 M urea Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. Store at -80°C after receiving vials. Storage: Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. NP 000236 RefSeq: 4233 Locus ID: UniProt ID: P08581, A0A024R759 **RefSeq Size:** 6641 Cytogenetics: 7q31.2 **RefSeq ORF:** 4170 Synonyms: AUTS9; c-Met; DFNB97; HGFR; RCCP2



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRIGENE MET (NM_000245) Human Recombinant Protein – TP762409

- Summary: This gene encodes a member of the receptor tyrosine kinase family of proteins and the product of the proto-oncogene MET. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that are linked via disulfide bonds to form the mature receptor. Further processing of the beta subunit results in the formation of the M10 peptide, which has been shown to reduce lung fibrosis. Binding of its ligand, hepatocyte growth factor, induces dimerization and activation of the receptor, which plays a role in cellular survival, embryogenesis, and cellular migration and invasion. Mutations in this gene are associated with papillary renal cell carcinoma, hepatocellular carcinoma, and various head and neck cancers. Amplification and overexpression of this gene are also associated with multiple human cancers. [provided by RefSeq, May 2016]
- Protein Families: Druggable Genome, Protein Kinase, Transmembrane
- Protein Pathways:Adherens junction, Axon guidance, Colorectal cancer, Cytokine-cytokine receptor interaction,
Endocytosis, Epithelial cell signaling in Helicobacter pylori infection, Focal adhesion,
Melanoma, Pathways in cancer, Renal cell carcinoma

Product images:



Purified recombinant protein MET was analyzed by SDS-PAGE gel and Coomossie Blue Staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US