

## **Product datasheet for TP790160**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## WNT3 (NM\_030753) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human wingless-type MMTV integration site family, member

3 (WNT3), with C-terminal DDK tag, secretory expressed in CHO cells, 20ug

Species: Human

**Expression Host:** CHO

Expression cDNA Clone

or AA Sequence:

A DNA sequence from TrueORF clone, RC211115, encoding the region Gly22-Lys355 of WNT3

Tag: C-DDK

Predicted MW: 37.4 kDa

**Concentration:** >0.05 μg/μL as determined by Bradford protein assay

**Purity:** > 90% as determined by SDS-PAGE and Coomassie blue staining

Buffer: PBS, pH 7.4, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 110380

 Locus ID:
 7473

 UniProt ID:
 P56703

RefSeq Size: 1506

**Cytogenetics:** 17q21.31-q21.32

RefSeq ORF: 1065

Synonyms: INT4; TETAMS





**Summary:** 

The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It encodes a protein which shows 98% amino acid identity to mouse Wnt3 protein, and 84% to human WNT3A protein, another WNT gene product. The mouse studies show the requirement of Wnt3 in primary axis formation in the mouse. Studies of the gene expression suggest that this gene may play a key role in some cases of human breast, rectal, lung, and gastric cancer through activation of the WNT-beta-catenin-TCF signaling pathway. This gene is clustered with WNT15, another family member, in the chromosome 17q21 region. [provided by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane

**Protein Pathways:** Basal cell carcinoma, Hedgehog signaling pathway, Melanogenesis, Pathways in cancer, Wnt

signaling pathway

## **Product images:**

