

## Product datasheet for **TR320083**

### Vasopressin (AVP) Human shRNA Plasmid Kit (Locus ID 551)

#### Product data:

Product Type:	shRNA Plasmids
Product Name:	Vasopressin (AVP) Human shRNA Plasmid Kit (Locus ID 551)
Locus ID:	551
Synonyms:	ADH; ARVP; AVP-NPII; AVRVP; VP
Vector:	pRS (TR20003)
E. coli Selection:	Ampicillin
Mammalian Cell Selection:	Puromycin
Format:	Retroviral plasmids
Components:	AVP - Human, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID = 551). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.
RefSeq:	<a href="#">NM_000490</a> , <a href="#">NM_000490.1</a> , <a href="#">NM_000490.2</a> , <a href="#">NM_000490.3</a> , <a href="#">NM_000490.4</a> , <a href="#">BC018062</a> , <a href="#">BC126196</a> , <a href="#">BC126224</a> , <a href="#">NM_000490.5</a>
UniProt ID:	<a href="#">P01185</a>



[View online »](#)

<b>Summary:</b>	<p>This gene encodes a member of the vasopressin/oxytocin family and preproprotein that is proteolytically processed to generate multiple protein products. These products include the neuropeptide hormone arginine vasopressin, and two other peptides, neurophysin 2 and copeptin. Arginine vasopressin is a posterior pituitary hormone that is synthesized in the supraoptic nucleus and paraventricular nucleus of the hypothalamus. Along with its carrier protein, neurophysin 2, it is packaged into neurosecretory vesicles and transported axonally to the nerve endings in the neurohypophysis where it is either stored or secreted into the bloodstream. The precursor is thought to be activated while it is being transported along the axon to the posterior pituitary. Arginine vasopressin acts as a growth factor by enhancing pH regulation through acid-base transport systems. It has a direct antidiuretic action on the kidney, and also causes vasoconstriction of the peripheral vessels. This hormone can contract smooth muscle during parturition and lactation. It is also involved in cognition, tolerance, adaptation and complex sexual and maternal behaviour, as well as in the regulation of water excretion and cardiovascular functions. Mutations in this gene cause autosomal dominant neurohypophyseal diabetes insipidus (ADNDI). This gene is present in a gene cluster with the related gene oxytocin on chromosome 20. [provided by RefSeq, Nov 2015]</p>
<b>shRNA Design:</b>	<p>These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. If you need a special design or shRNA sequence, please utilize our <a href="#">custom shRNA service</a>.</p>
<b>Performance Guaranteed:</b>	<p>OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.</p> <p>For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).</p>