

Product datasheet for **TR500696**

Fgfr2 Mouse shRNA Plasmid (Locus ID 14183)

Product data:

Product Type:	shRNA Plasmids
Product Name:	Fgfr2 Mouse shRNA Plasmid (Locus ID 14183)
Locus ID:	14183
Synonyms:	AU043015; AW556123; Bek; Fgfr-2; Fgfr-7; Fgfr7; KGFR; KGFRTr; svs
Vector:	pRS (TR20003)
E. coli Selection:	Ampicillin
Mammalian Cell Selection:	Puromycin
Format:	Retroviral plasmids
Components:	Fgfr2 - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID = 14183). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.
RefSeq:	BC091652 , NM_001347638 , NM_010207 , NM_201601 , NM_201601.1 , NM_201601.2 , NM_010207.1 , NM_010207.2 , BC065069 , BC072640 , BC151201 , BC172174
UniProt ID:	P21803



[View online »](#)

Summary:	<p>Tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of cell proliferation, differentiation, migration and apoptosis, and in the regulation of embryonic development. Required for normal embryonic patterning, trophoblast function, limb bud development, lung morphogenesis, osteogenesis and skin development. Plays an essential role in the regulation of osteoblast differentiation, proliferation and apoptosis, and is required for normal skeleton development. Promotes cell proliferation in keratinocytes and immature osteoblasts, but promotes apoptosis in differentiated osteoblasts. Phosphorylates PLCG1, FRS2 and PAK4. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. FGFR2 signaling is down-regulated by ubiquitination, internalization and degradation. Mutations that lead to constitutive kinase activation or impair normal FGFR2 maturation, internalization and degradation lead to aberrant signaling. Over-expressed FGFR2 promotes activation of STAT1.[UniProtKB/Swiss-Prot Function]</p>
shRNA Design:	<p>These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our custom shRNA service.</p>
Performance Guaranteed:	<p>OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.</p> <p>For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).</p>