

Product datasheet for TR501558

Slc22a6 Mouse shRNA Plasmid (Locus ID 18399)

Product data:

Product Type: shRNA Plasmids

Product Name: Slc22a6 Mouse shRNA Plasmid (Locus ID 18399)

Locus ID: 18399

Synonyms: mOat1; NKT; Oat1; Orctl1

Vector: pRS (TR20003)

E. coli Selection: Ampicillin

Mammalian Cell Puromycin

Selection:

Format: Retroviral plasmids

Components: Slc22a6 - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID =

18399). 5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.

RefSeq: <u>BC021647</u>, <u>NM 008766</u>, <u>NM 008766.1</u>, <u>NM 008766.2</u>, <u>NM 008766.3</u>

UniProt ID: Q8VC69

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com CN: techsupport@origene.cn



Summary:

Involved in the renal elimination of endogenous and exogenous organic anions. Functions as organic anion exchanger when the uptake of one molecule of organic anion is coupled with an efflux of one molecule of endogenous dicarboxylic acid (glutarate, ketoglutarate, etc). Mediates the sodium-independent uptake of 2,3-dimercapto-1-propanesulfonic acid (DMPS), cidofovir, adefovir, 9-(2-phosphonylmethoxyethyl) guanine (PMEG), 9-(2phosphonylmethoxyethyl) diaminopurine (PMEDAP), ochratoxin (OTA), acyclovir (ACV), 3'azido-3-'deoxythymidine (AZT), cimetidine (CMD), 2,4-dichloro-phenoxyacetate (2,4-D), hippurate (HA), indoleacetate (IA), indoxyl sulfate (IS) and 3-carboxy-4-methyl-5-propyl-2furanpropionate (CMPF) and edaravone sulfate (By similarity). Mediates the sodiumindependent uptake of p-aminohippurate (PAH). PAH uptake is inhibited by benzothiazolylcysteine (BTC), S-chlorotrifluoroethylcysteine (CTFC), cysteine S-conjugates Sdichlorovinylcysteine (DCVC), furosemide, steviol, phorbol 12-myristate 13-acetate (PMA), calcium ionophore A23187, benzylpenicillin, bumetamide, losartan, probenecid, phenol red, urate, glutarate and alpha-ketoglutarate (By similarity). PAH uptake is inhibited by pchloromercuribenzenesulphonate (PCMBS), diethyl pyrocarbonate (DEPC), indomethacin, sulindac, diclofenac, carprofen, okadaic acid and PKC activators.[UniProtKB/Swiss-Prot Function]

shRNA Design:

Performance Guaranteed: These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our custom shRNA service.

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).