

## **Product datasheet for TR501722**

## Ppp1ca Mouse shRNA Plasmid (Locus ID 19045)

**Product data:** 

**Product Type:** shRNA Plasmids

**Product Name:** Ppp1ca Mouse shRNA Plasmid (Locus ID 19045)

**Locus ID:** 19045

**Synonyms:** dism2; Ppp1c

Vector: pRS (TR20003)

E. coli Selection: Ampicillin

Mammalian Cell Puromycin

Selection:

•

Format: Retroviral plasmids

Components: Ppp1ca - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID =

19045). 5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.

RefSeq: <u>BC014828, NM 031868, NM 031868.1, NM 031868.2</u>

**UniProt ID:** <u>P62137</u>

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## Summary:

Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets. Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase. Regulates NEK2 function in terms of kinase activity and centrosome number and splitting, both in the presence and absence of radiation-induced DNA damage. Regulator of neural tube and optic fissure closure, and enteric neural crest cell (ENCCs) migration during development. In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. May dephosphorylate CSNK1D and CSNK1E. Dephosphorylates CENPA (By similarity). Dephosphorylates the 'Ser-139' residue of ATG16L1 causing dissociation of ATG12-ATG5-ATG16L1 complex, thereby inhibiting autophagy (By similarity).[UniProtKB/Swiss-Prot Function]

## shRNA Design:

Performance Guaranteed:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. If you need a special design or shRNA sequence, please utilize our <a href="mailto:custom shRNA service">custom shRNA service</a>.

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).