

Product datasheet for **TR502336**

Ctr9 Mouse shRNA Plasmid (Locus ID 22083)

Product data:

Product Type:	shRNA Plasmids
Product Name:	Ctr9 Mouse shRNA Plasmid (Locus ID 22083)
Locus ID:	22083
Synonyms:	AA409336; mKIAA0155; Sh2bp1; Tsbp; Tsp
Vector:	pRS (TR20003)
E. coli Selection:	Ampicillin
Mammalian Cell Selection:	Puromycin
Format:	Retroviral plasmids
Components:	Ctr9 - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID = 22083). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.
RefSeq:	NM_009431 , NM_009431.1 , NM_009431.2 , BC010344 , BC024749 , BC053910 , BC060091 , BC071247 , BC080719
UniProt ID:	Q62018



[View online »](#)

Summary:	<p>Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1); UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. Required for mono- and trimethylation on histone H3 'Lys-4' (H3K4me3) and dimethylation on histone H3 'Lys-79' (H3K4me3). Required for Hox gene transcription (By similarity). Required for the trimethylation of histone H3 'Lys-4' (H3K4me3) on genes involved in stem cell pluripotency; this function is synergistic with CXXC1 indicative for an involvement of the SET1 complex. Involved in transcriptional regulation of IL6-responsive genes and in JAK-STAT pathway; may regulate DNA-association of STAT3. [UniProtKB/Swiss-Prot Function]</p>
shRNA Design:	<p>These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our custom shRNA service.</p>
Performance Guaranteed:	<p>OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.</p> <p>For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).</p>