

# Product datasheet for TR502589

## Map2k1 Mouse shRNA Plasmid (Locus ID 26395)

## **Product data:**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type:                | shRNA Plasmids  |
|------------------------------|---|
| Product Name:                | Map2k1 Mouse shRNA Plasmid (Locus ID 26395)   |
| Locus ID:                    | 26395   |
| Synonyms:                    | MAPKK1; Mek1; MEKK1; Prkmk1   |
| Vector:                      | pRS (TR20003)   |
| E. coli Selection:           | Ampicillin  |
| Mammalian Cell<br>Selection: | Puromycin   |
| Format:                      | Retroviral plasmids   |
| Components:                  | Map2k1 - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID =<br>26395). 5µg purified plasmid DNA per construct<br>29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free. |
| RefSeq:                      | BC054754, NM 008927, NM 008927.1, NM 008927.2, NM 008927.3, BC051137, NM 008927.4   |
| UniProt ID:                  | <u>P31938</u>   |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** Map2k1 Mouse shRNA Plasmid (Locus ID 26395) – TR502589

Dual specificity protein kinase which acts as an essential component of the MAP kinase signal Summary: transduction pathway. Binding of extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1/MEK1 and MAP2K2/MEK2. Both MAP2K1/MEK1 and MAP2K2/MEK2 function specifically in the MAPK/ERK cascade, and catalyze the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in the extracellular signal-regulated kinases MAPK3/ERK1 and MAPK1/ERK2, leading to their activation and further transduction of the signal within the MAPK/ERK cascade. Depending on the cellular context, this pathway mediates diverse biological functions such as cell growth, adhesion, survival and differentiation, predominantly through the regulation of transcription, metabolism and cytoskeletal rearrangements. One target of the MAPK/ERK cascade is peroxisome proliferator-activated receptor gamma (PPARG), a nuclear receptor that promotes differentiation and apoptosis. MAP2K1/MEK1 has been shown to export PPARG from the nucleus. The MAPK/ERK cascade is also involved in the regulation of endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC), as well as in the fragmentation of the Golgi apparatus during mitosis.[UniProtKB/Swiss-Prot Function]

shRNA Design:These shRNA constructs were designed against multiple splice variants at this gene locus. To<br/>be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>.If you need a special design or shRNA sequence, please utilize our <a href="mailto:custom shRNA service">custom shRNA service</a>.

PerformanceOriGene guarantees that the sequences in the shRNA expression cassettes are verified toGuaranteed:correspond to the target gene with 100% identity. One of the four constructs at minimum are<br/>guaranteed to produce 70% or more gene expression knock-down provided a minimum<br/>transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to<br/>evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly<br/>assess knockdown, the gene expression level from the included scramble control vector must<br/>be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US