

Product datasheet for **TR508392**

Gpr183 Mouse shRNA Plasmid (Locus ID 321019)

Product data:

| | |
|---------------------------|---|
| Product Type: | shRNA Plasmids |
| Product Name: | Gpr183 Mouse shRNA Plasmid (Locus ID 321019) |
| Locus ID: | 321019 |
| Synonyms: | Ebi2 |
| Vector: | pRS (TR20003) |
| E. coli Selection: | Ampicillin |
| Mammalian Cell Selection: | Puromycin |
| Format: | Retroviral plasmids |
| Components: | Gpr183 - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID = 321019). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free. |
| RefSeq: | BC052868 , NM_183031 , NM_183031.1 , NM_183031.2 |
| UniProt ID: | Q3U6B2 |



[View online »](#)

Summary:

G-protein coupled receptor expressed in lymphocytes that acts as a chemotactic receptor for B-cells, T-cells, splenic dendritic cells, monocytes/macrophages and astrocytes (PubMed:19597478, PubMed:19615922, PubMed:21844396, PubMed:21796211, PubMed:21796212, PubMed:27147029). Receptor for oxysterol 7-alpha,25-dihydroxycholesterol (7-alpha,25-OHC) and other related oxysterols (PubMed:21796211, PubMed:21796212). Mediates cell positioning and movement of a number of cells by binding the 7-alpha,25-OHC ligand that forms a chemotactic gradient (PubMed:21796211, PubMed:21796212, PubMed:27147029). Binding of 7-alpha,25-OHC mediates the correct localization of B-cells during humoral immune responses (PubMed:21796211, PubMed:21796212). Collaborates with CXCR5 to mediate B-cell migration; probably by forming a heterodimer with CXCR5 that affects the interaction between of CXCL13 and CXCR5 (PubMed:21948984, PubMed:22913878). Guides B-cell movement along the B-cell zone-T-cell zone boundary and later to interfollicular and outer follicular regions (PubMed:19615922, PubMed:19597478, PubMed:21844396). Its specific expression during B-cell maturation helps position B-cells appropriately for mounting T-dependent antibody responses (PubMed:19615922). Also acts as a chemotactic receptor for some T-cells upon binding to 7-alpha,25-OHC ligand (PubMed:27147029). Promotes follicular helper T (Tfh) cells differentiation by positioning activated T-cells at the follicle-T-zone interface, promoting contact of newly activated CD4 T-cells with activated dendritic cells and exposing them to Tfh-cell-promoting inducible costimulator (ICOS) ligand (PubMed:27147029). Expression in splenic dendritic cells is required for their homeostasis, localization and ability to induce B- and T-cell responses: GPR183 acts as a chemotactic receptor in dendritic cells that mediates the accumulation of CD4(+) dendritic cells in bridging channels (PubMed:23682316, PubMed:23502855). Regulates migration of astrocytes and is involved in communication between astrocytes and macrophages (PubMed:25297897, PubMed:27166278). Promotes osteoclast precursor migration to bone surfaces (PubMed:26438360). Signals constitutively through G(i)-alpha, but not G(s)-alpha or G(q)-alpha (By similarity). Signals constitutively also via MAPK1/3 (ERK1/2) (By similarity).[UniProtKB/Swiss-Prot Function]

shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).

**Performance
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).