

Product datasheet for TR509872

Il1f6 Mouse shRNA Plasmid (Locus ID 54448)

Product data:

Product Type: shRNA Plasmids

Product Name: II1f6 Mouse shRNA Plasmid (Locus ID 54448)

Locus ID: 54448

Synonyms: Fil1; IL-1H1; Il1f9; IL1RP2; Il36a

Vector: pRS (TR20003)

E. coli Selection: Ampicillin

Mammalian Cell Puromycin

Selection:

Format: Retroviral plasmids

Components: Il1f6 - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID =

54448). 5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.

RefSeq: <u>BC117102</u>, <u>BC117104</u>, <u>NM 019450</u>, <u>NM 019450.1</u>, <u>NM 019450.2</u>, <u>NM 019450.3</u>

UniProt ID: Q9|LA2

Summary: Cytokine that binds to and signals through the IL1RL2/IL-36R receptor which in turn activates

NF-kappa-B and MAPK signaling pathways in target cells linked to a pro-inflammatory response. Part of the IL-36 signaling system that is thought to be present in epithelial barriers

and to take part in local inflammatory response; similar to the IL-1 system with which it shares the coreceptor IL1RAP. Seems to be involved in skin inflammatory response by acting on keratinocytes, dendritic cells and indirectly on T-cells to drive tissue infiltration, cell maturation and cell proliferation. Induces the production of proinflammatory cytokines, including IL-12, Il-1 beta, IL-6, TNF-alpha and IL-23 in bone marrow-derived dendritic cells (BMDCs). Involved in dendritic cell maturation by stimulating the surface expression of CD80, CD86 and MHC class II. Induces the production of IFN-gamma, IL-4 and IL-17 by cultured CD4(+) T-cells and splenocytes. May play a role in proinflammatory effects in the lung: induces the expression of CXCL1 and CXCL2 in the lung, and the expression of TNF-alpha, IL-36c, IL-1A, IL-1B, CXCL1 and CXCL2 in isolated splenic CD11c(+) alveolar macrophages. May be

Induces NF-kappa B activation in macrophages.[UniProtKB/Swiss-Prot Function]


involved in T-cell maturation by stimulating the surface expression of CD40 and modestly CD80 and CD86 in splenic CD11c(+) cells. May be involved in CD4(+) T-cell proliferation.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com





shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our custom shRNA service.

Performance Guaranteed: OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).