

## **Product datasheet for TR512428**

## **Rest Mouse shRNA Plasmid (Locus ID 19712)**

**Product data:** 

**Product Type:** shRNA Plasmids

**Product Name:** Rest Mouse shRNA Plasmid (Locus ID 19712)

**Locus ID:** 19712

**Synonyms:** 2610008J04Rik; AA407358; NRSF; REST4

**Vector:** pRS (TR20003)

E. coli Selection: Ampicillin

Mammalian Cell Puromycin

Selection:

Format: Retroviral plasmids

Components: Rest - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID =

19712). 5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.

**RefSeq:** <u>BC127065</u>, <u>NM 011263</u>, <u>NM 011263.1</u>, <u>NM 011263.2</u>, <u>BC023433</u>, <u>BC051648</u>, <u>BC096434</u>

UniProt ID: Q8VIG1

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Summary:** 

Transcriptional repressor which binds neuron-restrictive silencer element (NRSE) and represses neuronal gene transcription in non-neuronal cells (PubMed:29961578, PubMed:9771705). Restricts the expression of neuronal genes by associating with two distinct corepressors, SIN3A and RCOR1, which in turn recruit histone deacetylase to the promoters of REST-regulated genes (By similarity). Mediates repression by recruiting the BHC complex at RE1/NRSE sites which acts by deacetylating and demethylating specific sites on histones, thereby acting as a chromatin modifier (By similarity). Transcriptional repression by REST-CDYL via the recruitment of histone methyltransferase EHMT2 may be important in transformation suppression (By similarity). Represses the expression of SRRM4 in non-neural cells to prevent the activation of neural-specific splicing events and to prevent production of REST isoform 2 (PubMed:21884984). Repressor activity may be inhibited by forming heterodimers with isoform 2, thereby preventing binding to NRSE or binding to corepressors and leading to derepression of target genes (PubMed:10490617, PubMed:11039732). Also maintains repression of neuronal genes in neural stem cells, and allows transcription and differentiation into neurons by dissociation from RE1/NRSE sites of target genes (PubMed:15907476). Thereby is involved in maintaining the quiescent state of adult neural stem cells and preventing premature differentiation into mature neurons (PubMed:27819263). Plays a role in the developmental switch in synaptic NMDA receptor composition during postnatal development, by repressing GRIN2B expression and thereby altering NMDA receptor properties from containing primarily GRIN2B to primarily GRIN2A subunits (By similarity). Acts as a regulator of osteoblast differentiation (PubMed:25727884). Key repressor of gene expression in hypoxia; represses genes in hypoxia by direct binding to an RE1/NRSE site on their promoter regions (By similarity). May also function in stress resistance in the brain during aging; possibly by regulating expression of genes involved in cell death and in the stress response (PubMed:24670762). Repressor of gene expression in the hippocampus after ischemia by directly binding to RE1/NRSE sites and recruiting SIN3A and RCOR1 to promoters of target genes, thereby promoting changes in chromatin modifications and ischemia-induced cell death (By similarity). After ischemia, might play a role in repression of miR-132 expression in hippocampal neurons, thereby leading to neuronal cell death (By similarity).[UniProtKB/Swiss-Prot Function]

shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact <a href="mailto:techsupport@origene.com">techsupport@origene.com</a>. If you need a special design or shRNA sequence, please utilize our <a href="mailto:custom shRNA service">custom shRNA service</a>.



## Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).