

Product datasheet for **TR515640**

Blk Mouse shRNA Plasmid (Locus ID 12143)

Product data:

Product Type:	shRNA Plasmids
Product Name:	Blk Mouse shRNA Plasmid (Locus ID 12143)
Locus ID:	12143
Vector:	pRS (TR20003)
E. coli Selection:	Ampicillin
Mammalian Cell Selection:	Puromycin
Format:	Retroviral plasmids
Components:	Blk - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID = 12143). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.
RefSeq:	BC030668 , NM_007549 , NM_007549.1 , NM_007549.2
UniProt ID:	P16277



[View online »](#)

Summary:	<p>Non-receptor tyrosine kinase involved in B-lymphocyte development, differentiation and signaling. B-cell receptor (BCR) signaling requires a tight regulation of several protein tyrosine kinases and phosphatases, and associated coreceptors (PubMed:2404338, PubMed:7690139, PubMed:7608542, PubMed:9636152, PubMed:14662906, PubMed:12563261). Binding of antigen to the B-cell antigen receptor (BCR) triggers signaling that ultimately leads to B-cell activation (PubMed:2404338, PubMed:7690139, PubMed:7608542, PubMed:14662906, PubMed:12563261). Signaling through BLK plays an important role in transmitting signals through surface immunoglobulins and supports the pro-B to pre-B transition, as well as the signaling for growth arrest and apoptosis downstream of B-cell receptor (PubMed:2404338, PubMed:7690139, PubMed:7608542, PubMed:14662906, PubMed:12563261). Specifically binds and phosphorylates CD79A at 'Tyr-188' and 'Tyr-199', as well as CD79B at 'Tyr-196' and 'Tyr-207' (PubMed:7592958, PubMed:9177269). Phosphorylates also the immunoglobulin G receptor FCGR2 (By similarity). With FYN and LYN, plays an essential role in pre-B-cell receptor (pre-BCR)-mediated NF-kappa-B activation (PubMed:14662906, PubMed:12563261). Contributes also to BTK activation by indirectly stimulating BTK intramolecular autophosphorylation (PubMed:7565679). In pancreatic islets, acts as a modulator of beta-cells function through the up-regulation of PDX1 and NKX6-1 and consequent stimulation of insulin secretion in response to glucose (By similarity). Phosphorylates CGAS, promoting retention of CGAS in the cytosol (By similarity).[UniProtKB/Swiss-Prot Function]</p>
shRNA Design:	<p>These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our custom shRNA service.</p>
Performance Guaranteed:	<p>OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.</p> <p>For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).</p>