

Product datasheet for TR515725

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Dclk1 Mouse shRNA Plasmid (Locus ID 13175)

Product data:

Product Type: shRNA Plasmids

Product Name: Dclk1 Mouse shRNA Plasmid (Locus ID 13175)

Locus ID: 13175

Synonyms: 1700113D08Rik; 2810480F11Rik; Al836758; Clic; Click-I; CPG1; Cpg16; Dc; Dcamk; Dcamkl1;

Dcl; Dclk; mKIAA0369

Vector: pRS (TR20003)

E. coli Selection: Ampicillin

Mammalian Cell Puromycin

Selection:

_

Format: Retroviral plasmids

Components: Dclk1 - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID =

13175). 5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.

RefSeq: <u>BC133685, NM 001111051, NM 001111052, NM 001111053, NM 001195538, NM 001195539</u>,

NM 001195540, NM 019978, NM 019978.1, NM 019978.2, NM 019978.3, NM 001111051.1, NM 001111052.1, NM 001111053.1, NM 001195538.1, NM 001195539.1, NM 001195540.1,

BC021354, BC050903, BC064783, BM933369, NM 001111052.2, NM 001111053.2,

NM 001111051.2, NM 019978.4, NM 001195540.2, NM 001195539.2

UniProt ID: Q9|LM8



Summary:

This gene encodes a member of the protein kinase superfamily and the doublecortin family. The protein encoded by this gene contains two N-terminal doublecortin domains, which bind microtubules and regulate microtubule polymerization, a C-terminal serine/threonine protein kinase domain, which shows substantial homology to Ca2+/calmodulin-dependent protein kinase, and a serine/proline-rich domain in between the doublecortin and the protein kinase domains, which mediates multiple protein-protein interactions. The microtubule-polymerizing activity of the encoded protein is independent of its protein kinase activity. The encoded protein is involved in several different cellular processes, including neuronal migration, retrograde transport, neuronal apoptosis and neurogenesis. This gene is upregulated by brain-derived neurotrophic factor and associated with memory and general cognitive abilities. Multiple transcript variants generated by two alternative promoter usage and alternative splicing have been found, but the biological validity of some variants has not been determined. These variants encode different isoforms, which are differentially expressed and have different kinase activities. [provided by RefSeq, Sep 2010]

shRNA Design:

Performance Guaranteed: These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our custom shRNA service.

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).