

Product datasheet for **TR518976**

Itch Mouse shRNA Plasmid (Locus ID 16396)

Product data:

Product Type:	shRNA Plasmids
Product Name:	Itch Mouse shRNA Plasmid (Locus ID 16396)
Locus ID:	16396
Synonyms:	6720481N21Rik; 8030492O04Rik; A130065M08; AIP4; C230047C07Rik
Vector:	pRS (TR20003)
E. coli Selection:	Ampicillin
Mammalian Cell Selection:	Puromycin
Format:	Retroviral plasmids
Components:	Itch - Mouse, 4 unique 29mer shRNA constructs in retroviral untagged vector(Gene ID = 16396). 5µg purified plasmid DNA per construct 29-mer scrambled shRNA cassette in pRS Vector, TR30012, included for free.
RefSeq:	BC062934 , BC064678 , NM_001243712 , NM_008395 , NR_037196 , NM_008395.1 , NM_008395.2 , NM_008395.3 , NM_001243712.1 , BC048489
UniProt ID:	Q8C863



[View online »](#)

Summary:

Acts as an E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates (PubMed:15358865, PubMed:16446428, PubMed:17592138, PubMed:18628966, PubMed:20392206, PubMed:25632008). It catalyzes 'Lys-29-', 'Lys-48'- and 'Lys-63'-linked ubiquitin conjugation (By similarity). Involved in the control of inflammatory signaling pathways (By similarity). Is an essential component of a ubiquitin-editing protein complex, comprising also TNFAIP3, TAX1BP1 and RNF11, that ensures the transient nature of inflammatory signaling pathways (By similarity). Promotes the association of the complex after TNF stimulation (By similarity). Once the complex is formed, TNFAIP3 deubiquitinates 'Lys-63' polyubiquitin chains on RIPK1 and catalyzes the formation of 'Lys-48'-polyubiquitin chains (By similarity). This leads to RIPK1 proteasomal degradation and consequently termination of the TNF- or LPS-mediated activation of NFkB1 (By similarity). Ubiquitinates RIPK2 by 'Lys-63'-linked conjugation and influences NOD2-dependent signal transduction pathways (By similarity). Regulates the transcriptional activity of several transcription factors involved in immune response (PubMed:15358865, PubMed:11828324). Ubiquitinates NFE2 by 'Lys-63' linkages and is implicated in the control of the development of hematopoietic lineages (By similarity). Mediates JUN ubiquitination and degradation (PubMed:15358865). Mediates JUNB ubiquitination and degradation (PubMed:11828324, PubMed:15358865). Critical regulator of type 2 helper T (Th2) cell cytokine production by inducing JUNB ubiquitination and degradation (PubMed:11828324). Involved in the negative regulation of MAVS-dependent cellular antiviral responses (By similarity). Ubiquitinates MAVS through 'Lys-48'-linked conjugation resulting in MAVS proteasomal degradation (By similarity). Following ligand stimulation, regulates sorting of Wnt receptor FZD4 to the degradative endocytic pathway probably by modulating PI42KA activity (By similarity). Ubiquitinates PI4K2A and negatively regulates its catalytic activity (By similarity). Ubiquitinates chemokine receptor CXCR4 and regulates sorting of CXCR4 to the degradative endocytic pathway following ligand stimulation by ubiquitinating endosomal sorting complex required for transport ESCRT-0 components HGS and STAM (By similarity). Targets DTX1 for lysosomal degradation and controls NOTCH1 degradation, in the absence of ligand, through 'Lys-29'-linked polyubiquitination (PubMed:18628966). Ubiquitinates SNX9 (By similarity). Ubiquitinates MAP3K7 through 'Lys-48'-linked conjugation (PubMed:25632008). Involved in the regulation of apoptosis and reactive oxygen species levels through the ubiquitination and proteasomal degradation of TXNIP (By similarity). Mediates the antiapoptotic activity of epidermal growth factor through the ubiquitination and proteasomal degradation of p15 BID (PubMed:20392206). Ubiquitinates BRAT1 and this ubiquitination is enhanced in the presence of NDFIP1 (By similarity).[UniProtKB/Swiss-Prot Function]

shRNA Design:

These shRNA constructs were designed against multiple splice variants at this gene locus. To be certain that your variant of interest is targeted, please contact techsupport@origene.com. If you need a special design or shRNA sequence, please utilize our [custom shRNA service](#).

**Performance
Guaranteed:**

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).