

Product datasheet for TS402471P5

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MAGEA11 CytoSection

Product data:

Product Type: CytoSections

Description: Transient overexpression of MAGEA11, transcript variant 1, in HEK293T cells, FFPE control for

IHC, ICC and ISH staining, 25 slides per pack

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

TrueORF Clone RC202471

Tag: C-MYC/DDK

Detection Antibodies: DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)

ACCN: <u>NM 005366</u>, <u>NP 005357</u>

Synonyms: CT1.11; MAGE-11; MAGE11; MAGEA-11

Storage: Room Temperature

Stability: Slides are guaranteed for a year from the date of receipt if proper storage instructions were

followed.

Preparation: HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells

were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin &

dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut

and mounted on positively charged SuperFrost slides.

Note: This product is for research use only and is not approved for use in humans or in clinical


diagnosis.

RefSeq: NP 005357


Locus ID: 4110 Cytogenetics: Xq28



Product images:



Immunohistochemistry staining on MAGEA11 overexpressed cytosection TS402471P5 with mouse anti DDK clone OTI11C3 C/N [TA180144] at 1:600 dilution 1hr room temperature. Antibody staining was achieved with HIER Citrate pH6, Polink1 with DAB chromogen detection kit ([D12-18]). Positive stain shown with the brown chromogen present. mammalian cells transfected with cDNA clone [RC202471], 5 micron sections, 40x magnification.



Immunohistochemistry staining on MAGEA11 overexpressed cytosection TS402471P5 with mouse anti DDK clone OTI11C3 C/N [TA180144] at 1:600 dilution 1hr room temperature. Antibody staining was achieved with HIER Citrate pH6, Polink1 with DAB chromogen detection kit ([D12-18]). Positive stain shown with the brown chromogen present. mammalian cells transfected with cDNA clone [RC202471], 5 micron sections, 40x magnification.