

## Product datasheet for **TS404482P5**


### MAGEA4 CytoSection

#### Product data:


Product Type:	CytoSections
Description:	Transient overexpression of MAGEA4, transcript variant 3, in HEK293T cells, FFPE control for IHC, ICC and ISH staining, 25 slides per pack
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	TrueORF Clone RC204482
Tag:	C-MYC/DDK
Detection Antibodies:	DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)
Target Detection Antibodies:	MAGEA4 Mouse Monoclonal Antibody [Clone ID: OT11F9] (TA505362)
ACCN:	<a href="#">NM_001011549</a> , <a href="#">NP_001011549</a>
Synonyms:	CT1.4; MAGE-41; MAGE-X2; MAGE4; MAGE4A; MAGE4B
Storage:	Room Temperature
Stability:	Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.
Preparation:	HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.
Note:	This product is for research use only and is not approved for use in humans or in clinical diagnosis.
RefSeq:	<a href="#">NP_001011549</a>
Locus ID:	4103
Cytogenetics:	Xq28



[View online »](#)

**Product images:**

Immunohistochemistry staining on MAGEA4 variant 3 overexpressed cytosection TS404482P5 with mouse anti DDK clone OTI11C3 C/N [TA180144] at 1:600 dilution 1hr room temperature. Antibody staining was achieved with HIER Citrate pH6 , Polink1 with DAB chromogen detection kit ([D12-18]). Positive stain shown with the brown chromogen present. mammalian cells transfected with cDNA clone [RC204482], 5 micron sections, 40x magnification.



Immunohistochemistry staining on MAGEA4 variant 3 overexpressed cytosection TS404482P5 with mouse anti DDK clone OTI11C3 C/N [TA180144] at 1:600 dilution 1hr room temperature. Antibody staining was achieved with HIER Citrate pH6 , Polink1 with DAB chromogen detection kit ([D12-18]). Positive stain shown with the brown chromogen present. mammalian cells transfected with cDNA clone [RC204482], 5 micron sections, 40x magnification.