

## **Product datasheet for TS404894P5**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Fas Ligand (FASLG) CytoSection

**Product data:** 

**Product Type:** CytoSections

**Description:** Transient overexpression of FASLG in HEK293T cells, FFPE control for IHC, ICC and ISH

staining, 25 slides per pack

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** 

or AA Sequence:

TrueORF Clone RC204894

Tag: C-MYC/DDK

**Detection Antibodies:** DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)

ACCN: <u>NM 000639</u>, <u>NP 000630</u>

Synonyms: ALPS1B; APT1LG1; APTL; CD95-L; CD95L; CD178; FASL; TNFSF6; TNLG1A

Storage: Room Temperature

**Stability:** Slides are guaranteed for a year from the date of receipt if proper storage instructions were

followed.

**Preparation:** HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells

were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin &

dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut

and mounted on positively charged SuperFrost slides.

**Note:** This product is for research use only and is not approved for use in humans or in clinical

diagnosis.

RefSeq: NP 000630

Locus ID: 356

Cytogenetics: 1q24.3

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane


**Protein Pathways:** Allograft rejection, Apoptosis, Autoimmune thyroid disease, Cytokine-cytokine receptor

interaction, Graft-versus-host disease, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Pathways in cancer, Type I diabetes mellitus





## **Product images:**



Immunohistology staining on FASLG overexpressed cytosection TS404894P5 with mouse anti DDK clone OTI11C3 C/N [TA180144] at 1:400 dilution 4C O/N. Antibody staining was achieved with HIER Citrate pH6 , Polink1 with DAB chromogen detection kit ([D11-18]). Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC204894], 5 micron sections, 40x magnification