

## Product datasheet for TS408975P5

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ICOS Ligand (ICOSLG) CytoSection

**Product data:** 

**Product Type:** CytoSections

**Description:** Transient overexpression of ICOSLG in HEK293T cells, FFPE control for IHC, ICC and ISH

staining, 25 slides per pack

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

TrueORF Clone RC208975

Tag: C-MYC/DDK

**Detection Antibodies:** DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)

**Target Detection** 

**Antibodies:** 

ICOS Ligand (ICOSLG) Mouse Monoclonal Antibody [Clone ID: OTI2C5] (TA808779)

ACCN: <u>NM 015259, NP 056074</u>

**Synonyms:** B7-H2; B7h; B7H2; B7RP-1; B7RP1; CD275; GL50; ICOS-L; ICOSL; LICOS

Storage: Room Temperature

**Stability:** Slides are guaranteed for a year from the date of receipt if proper storage instructions were

followed.

**Preparation:** HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells

were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin &

dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut

and mounted on positively charged SuperFrost slides.

**Note:** This product is for research use only and is not approved for use in humans or in clinical

diagnosis.

**RefSeq:** NP 056074

**Locus ID:** 23308 **Cytogenetics:** 21q22.3


**Protein Families:** Transmembrane

**Protein Pathways:** Cell adhesion molecules (CAMs)





## **Product images:**



Immunohistology staining on ICOSLG overexpressed cytosection TS408975P5 with rabbit anti DDK clone OTIR5G2 C/N [TA592569] at 1:2000 dilution 20m RT. Antibody staining was achieved with HIER Citrate pH6 , Polink1 with DAB chromogen detection kit ([D11-18]). Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC208975], 5 micron sections, 40x magnification