

## **Product datasheet for TS409993P5**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Interferon gamma (IFNG) CytoSection

**Product data:** 

**Product Type:** CytoSections

**Description:** Transient overexpression of IFNG in HEK293T cells, FFPE control for IHC, ICC and ISH staining,

25 slides per pack

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

TrueORF Clone RC209993

Tag: C-MYC/DDK

**Detection Antibodies:** DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)

**Target Detection** 

Antibodies:

Interferon gamma (IFNG) Mouse Monoclonal Antibody [Clone ID: OTI1G7] (TA805797)

ACCN: <u>NM 000619, NP 000610</u>

Synonyms: IFG; IFI; IMD69

Storage: Room Temperature

**Stability:** Slides are guaranteed for a year from the date of receipt if proper storage instructions were

followed.

**Preparation:** HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells

were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin &

dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut

and mounted on positively charged SuperFrost slides.

**Note:** This product is for research use only and is not approved for use in humans or in clinical

diagnosis.

RefSeq: NP 000610

Locus ID: 3458 Cytogenetics: 12q15

**Protein Families:** Druggable Genome, Secreted Protein






**Protein Pathways:** 

Allograft rejection, Cytokine-cytokine receptor interaction, Graft-versus-host disease, Jak-STAT signaling pathway, Natural killer cell mediated cytotoxicity, Proteasome, Regulation of autophagy, Systemic lupus erythematosus, T cell receptor signaling pathway, TGF-beta signaling pathway, Type I diabetes mellitus

## **Product images:**



Immunohistology staining on IFNG overexpressed cytosection TS409993P5 with rabbit anti DDK clone OTIR5G2 C/N [TA592569] at 1:2000 dilution 20m RT. Antibody staining was achieved with HIER Citrate pH6 , Polink1 with DAB chromogen detection kit ([D11-18]). Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC209993], 5 micron sections, 40x magnification